

RAFDONAC® GEL

Diclofenac Diethylamine I.P. 1.16% w/w (eq. to Diclofenac Sodium) + Linseed Oil 1% w/w + Methyl Salicylate 10% w/w + Menthol 5% w/w

"Trusted Relief - Pain, Swelling, Stiffness, Mobility"

Marketed by: Human Orthocare

A Subsidiary of Human Biolife India Pvt. Ltd.

www.humanorthocare.com





Product Overview



Composition (per 30 g pack):

Diclofenac Diethylamine I.P.

1.16% w/w (eq. to Diclofenac Sodium)

Linseed Oil

1% w/w

Methyl Salicylate I.P.

10% w/w


Menthol I.P.

5% w/w

Key Features:

- Clinically proven topical NSAID
- Combines anti-inflammatory + counterirritant + cooling action
- Provides fast, localised pain relief
- Non-greasy & quick absorbing

Mechanism of Action



Diclofenac Diethylamine

Inhibits COX → reduces prostaglandins → ↓ inflammation & pain



Linseed Oil

Anti-inflammatory, improves penetration of actives



Methyl Salicylate

Counterirritant → improves blood circulation, reduces stiffness



Menthol


Cooling action → soothing comfort & faster relief



Result → Relieves pain, reduces swelling, restores mobility









Indications

Rafdonac® Gel is recommended for:

Arthritis Pain

Osteoarthritis & rheumatoid arthritis pain

Spinal Conditions

Low back pain & cervical spondylosis

Muscle Injuries

Musculoskeletal sprains & strains

Sports Medicine

Sports injuries & tendonitis


Trauma Care

Post-traumatic joint & muscle pain

Clinical Advantages









Localised Action

Localised action with minimal systemic exposure



Safety Profile

Safer alternative vs oral NSAIDs in high-risk patients



Patient-Friendly

Easy application, quick drying, non-greasy

Reduces pain, stiffness & swelling - Effective in ortho, trauma & sports injury practice



Comparison with Oral NSAIDs

Parameter

Localised Action

Systemic Side Effects

Onset of Relief

GI Safety

Long-Term Suitability

Rafdonac® Gel (Topical)

High

Minimal

Quick

Excellent

Safe

Oral NSAIDs

X Low

X High


✓ Systemic

X Poor

1 Limited



Dosage & Administration





01

Apply 2-4 g gel

To affected area 3–4 times daily

02

Massage gently

Until fully absorbed

- Avoid broken skin, eyes & mucous membranes
 - Wash hands after use

Safety & Tolerability



Generally Well-Tolerated

Possible mild side effects:

• Local redness, itching, burning sensation (rare)



Contraindications

- Hypersensitivity to diclofenac, salicylates or menthol
- Not suitable for children <12 years
- Avoid prolonged use in pregnancy/lactation







Why Prescribe Rafdonac® Gel?

- **3** Doctor's Choice For:
- Proven Efficacy

 Proven topical NSAID gel for musculoskeletal pain
- Triple Action

 Combines pain relief + anti-inflammatory + soothing effect
- Long-term Safety

 Safer long-term use vs oral NSAIDs





Contact Us

Step into care that moves you forward.

Use with clinical discretion, especially in high-risk orthopedic conditions.

HumAn Orthocare – A Subsidiary of Human Biolife India Pvt. Ltd.

For medical queries or collaboration:



www.humanorthocare.com